

For each of the following situations (1 and 2):

(A) Define variables in a tabular format, as follows.

name                      symbol                      scale

scale = nominal, ordinal, or cardinal  
cardinal = interval or ratio scale.

nv = number of variables  
nt = number of terms  
A. score = 3nv  
B. score = nt  
C. score = 2nv + 2  
D. score = 1

(B) Using the symbols, write a general linear model relating the response variable to explanatory variable(s) and interaction terms (if appropriate).

(C) Complete the first two columns of the ANOVA table                      source   df

(D) State the name of the analysis, from the following list.  
t-test, one-way ANOVA, two-way ANOVA, three-way ANOVA  
paired comparisons, randomized blocks,  
hierarchical (nested) ANOVA  
regression, multiple regression,  
ANCOVA (at least 1 nominal and at least 1 cardinal scale explanatory variable)  
none of the above.

1. Height is frequently named as a good predictor variable of weight among people of the same age and gender. Roberts (*American Journal of Clinical Nutrition* 54:499) measured the heights (cm) and weights (kg) of 14 males between the ages of 19 and 26 years of age. Does weight depend on height ?  
A=6 B=3 C=6 D=1

A. name   symbol   scale

C. source   df

B. \_\_\_\_\_ = \_\_\_\_\_ +  $\epsilon$    [3]

D. \_\_\_\_\_ [1]

2. Skinner and Allison (*J. Agric. Res.* 23:433-445) studied the effect of date of planting and amount of fertilizer (borax) on cotton growth, measured in pounds. Amount of borax was 0, 5, or 10 pounds. Three methods of borax application were (borax in drill & seed planted immediately, borax in drill & seed planted one week later, or borax broadcast). The experiment was carried out on 3 dates. When the analysis is carried out, all of the interaction terms were found to be non significant, with p-values of 0.173 or more. Write the model with no interaction terms.

A=12 B=5 C=10 D=1

A. <u>name</u> <u>symbol</u> <u>scale</u>
---

C. <u>source</u> <u>df</u>
----------------------------

$$\text{_____} = \text{_____} + \epsilon \quad [3]$$

D. [1]

3a. Define a symbol for scutum width (units of  $\mu\text{m}$ ) of ticks on rabbit #1 (Sokal and Rohlf, 1995, p 210), then define a symbol for the observed (sample) mean and the true (population) mean ..... [3]

3b. For the data on scutum width (8 values below) write the observed mean.  $\frac{\text{Symbol}}{\text{(Symbol)}} = \frac{\text{Value}}{\text{(Value)}} [1]$

3c. Write a probability statement for the 95% confidence limits around the true mean ..... [2]

3d. What value of the t-distribution should you use for the 95% limits ? \_\_\_\_\_ [1]

```
MTB > invcdf c1; SUBC> t 7.
0.0100 -2.9980
0.0250 -2.3646
0.0500 -1.8946
0.1000 -1.4149
0.9000 1.4149
0.9500 1.8946
0.9750 2.3646
0.9900 2.9980
```

3e. Compute the 95% confidence limits ..... [2]

```
MTB > print c2
ScWidth 380 376 360 368 372 366 374 382

MTB > describe c2
ScWidth N MEAN MEDIAN TRMEAN STDEV SEMEAN
8 372.25 373.00 372.25 7.36 2.60
```

4a. Construct an ANOVA table for which the total Sum of Squares is 100, 15% of this variability is due to regression, and the sample size is 10. Be sure to compute MS and F-ratio ..... [12]

4b. Explain how you would compute a p-value for the F-ratio in the table you have constructed, if the residuals were heterogeneous and non normal ..... [2]

4c. Circle the effect (increase/decrease) of doubling the sample size, in the ANOVA table you constructed (or any ANOVA table for regression) ..... [3]

increase decrease                      in MS error

increase decrease                      in F-ratio

increase decrease                      in p-value

For each of the following situations (1 and 2):

(A) Define variables in a tabular format, as follows.

name                      symbol                      scale

scale = nominal, ordinal, or cardinal  
cardinal = interval or ratio scale.

nv = number of variables  
nt = number of terms  
A. score = 3nv  
B. score = nt  
C. score = 2nv + 2  
D. score = 1

(B) Using the symbols, write a general linear model relating the response variable to explanatory variable(s) and interaction terms (if appropriate).

(C) Complete the first two columns of the ANOVA table                      source   df

(D) State the name of the analysis, from the following list.  
t-test, one-way ANOVA, two-way ANOVA, three-way ANOVA  
paired comparisons, randomized blocks,  
hierarchical (nested) ANOVA  
regression, multiple regression,  
ANCOVA (at least 1 nominal and at least 1 cardinal scale explanatory variable)  
none of the above.

1. Daniel (*Biostatistics* 1995, p234) reported cell diameters ( $\mu\text{m}$ ) of 40 lymphocytes and 50 tumor cells obtained from biopsies of tissue from patients with melanoma. Do cancerous and non-cancerous cells differ in diameter?                      A=6 B=3 C=6 D=1

A. name   symbol   scale

C. source   df

B. \_\_\_\_\_ = \_\_\_\_\_ +  $\epsilon$                       [3]

D. \_\_\_\_\_                      [1]

2. Does birth weight depend on maternal smoking, controlled for gestation period and maternal weight? Selvin (*Practical Biostatistical Methods*, 1995, Duxbury Press) reported birth weights of first infants (grams), gestation period (weeks), maternal smoking (0, 10-20, or  $\geq 40$  cigarettes per day), and maternal weight (kg) for 48 women over 40 years old. (Assume no interactive effects of explanatory variables on the response variable, as in multiple regression). A=12 B=5 C=10 D=1

A. name   symbol   scale

C. source   df

B. \_\_\_\_\_ = \_\_\_\_\_ +  $\epsilon$    [3]

D. \_\_\_\_\_ [1]

3a. Construct a one-way ANOVA table for which the total Sum of Squares is 100, 15% of this variability is due to treatment effects (control vs one treatment), and the sample size is 100. Be sure to compute MS and F-ratio ..... [12]

3b. Explain how you would compute a p-value for the F-ratio in the table you have constructed, if the residuals were normal and independent, with fixed variance ..... [2]

3c. Circle the effect (increase/decrease) of halving the sample size, in the ANOVA table you constructed (or any ANOVA table with the same model structure) ..... [3]

increase decrease                      in MS error

increase decrease                      in F-ratio

increase decrease                      in p-value

4a. Define a symbol for scutum width, in microns, of tick larvae on rabbit #4 (Sokal and Rohlf, 1995, p 208), then define a symbol for the observed (sample) mean and the true (population) mean ..... [3]

4b. For the data on scutum width (6 values below) write the observed mean.  $\frac{\text{(Symbol)}}{\text{(Value)}} = \frac{\text{[1]}}{\text{[1]}}$

4c. Write a probability statement for the 95% confidence limits around the true mean ..... [2]

4d. What value of the t-distribution should you use for the 95% limits ? \_\_\_\_\_ [1]

```
MTB > invcdf c10; SUBC> t 5.
0.0100 -3.3649
0.0250 -2.5706
0.0500 -2.0151
0.1000 -1.4759
0.9000 1.4759
0.9500 2.0151
0.9750 2.5706
0.9900 3.3649
```

4e. Compute the 95% confidence limits ..... [2]

```
C6
376 344 342 372 374 360
```

```
MTB > describe c6
```

		N	MEAN	MEDIAN	TRMEAN	STDEV	SEMEAN
C6	ScWidth	6	361.33	366.00	361.33	15.27	6.23